

**DEVELOPMENT OF LIMITING DILUTION VIABILITY PCR METHOD
TO ASSESS THE EFFECTIVENESS OF SELECTED BIOCIDES TO TREAT
INDOOR FUNGI GROWTH**

ER CHIN MING

A thesis submitted in
fulfillment of the requirement for the award of the
Doctor of Philosophy



Faculty of Civil and Environmental Engineering
Universiti Tun Hussein Onn Malaysia

JANUARY 2019

ACKNOWLEDGEMENT

First of all, I would like to express my sincere appreciation and gratitude to my bright and helpful supervisor, Associate Professor Ts. Dr. Norshuhaila binti Mohamed Sunar, co-supervisors, Associate Professor Engr. Dr. Abdul Mutalib bin Leman and Associate Professor Dr. Norzila binti Othman for giving me precious guidance, opinions and comments throughout study. I registered as a part time PhD candidate in the Faculty of Civil and Environmental Engineering, Universiti Tun Hussein Onn Malaysia on 3rd December 2012. The first few months of this study, I had to travel from my previous working place, Kota Tinggi to Parit Raja, Batu Pahat every weekend in order to carry out my experiments. Therefore, special thanks again to all my supervisor and co-supervisor for their understandings and helps. Other than that, special credits to all laboratory officers and assistants and other postgraduate students for all priceless advices, explanations and suggestions.

Besides, I am thankful for myself for being determined in this pursuit of great knowledge and of self-satisfaction. Being a part-time student, I had to expend my after-working-hours in this study. Determination in terms of physical and emotional was the power that pushed me throughout the study.

Finally, a special dedication to my family especially my wife, Madam Lee Chee Moi, who has been supporting and encouraging me to further my studies as a Doctor of Philosophy (PhD) candidate. Her support was magnificent and held me on the way of completing my PhD study. My mother, Madam Teo Suat and my father, Mr Er Swee Chai supported me to complete my thesis writing as soon as possible. Their support have provided me strength and courage and driven me throughout the study.

ABSTRACT

Indoor fungal contamination should be treated with cost-effective and green methods. Biocides have direct biological effect on living organisms but the evidence on their control of indoor fungal contamination is scarce. Using conventional cultivation to evaluate their effectiveness is time consuming while polymerase chain reaction (PCR) provides a fast and reliable alternative. The incorporation of serial dilution technique and viability information in PCR has made it suitable to evaluate the effectiveness of biocides. Thus, this study aimed to assess the antifungal ability of biocides, zinc salicylate (ZS), calcium benzoate (CB) and potassium sorbate (KS) to treat indoor fungal contamination through developing limiting dilution viability PCR (vPCR). These biocides were selected as they successfully controlled the growth of indoor waterborne fungi previously. Indoor air sampling revealed that higher educational building of computer studies (Building A) and of civil engineering studies (Building B) were contaminated by 509 CFU/m³ and 805.7 CFU/m³ of indoor airborne fungi, respectively. Two indoor fungi, *Talaromyces* spp. and *Aspergillus niger* were identified. They were subjected to biocides-treatment and subsequent conventional cultivation and limiting dilution vPCR due to their potential risks against humans' health. The limiting dilution vPCR was developed by incorporating the pre-treatment of propidium monoazide (PMA) before deoxyribonucleic acid (DNA) extraction and the serial dilution of the DNA template in PCR. This approach was proven to effectively enumerate the effectiveness of biocides to treat indoor fungi. KS was shown to have the best effectiveness (100%) to prevent the growth of *Talaromyces* spp, followed by ZS (80.8%) and CB (no effect). KS also showed the best effectiveness against *A. niger* (100%) at the early stage of the study but its effect reduced with time. ZS showed durable effect (66.67%) against *A. niger* Day 9 cultures. Inconstant results were indicated by cultivation method. This study has provided a cheaper, more accurate and suitable approach to determine the effectiveness of treatment of indoor fungi than cultivation methods.

ABSTRAK

Pencemaran kulat dalaman perlu dirawat dengan kaedah kos efektif dan mesra alam. Biosida berkesan langsung terhadap organisma tetapi bukti mereka dalam pemulihan pencemaran kulat dalaman berkekurangan. Penilaian kecekapan mereka dengan teknik pengkulturan konvensional sangat mengambil masa. Reaksi rantai polimerase (PCR) menyediakan alternatif yang cepat dan boleh dipercayai. Penggabungan teknik pencairan bersiri dan maklumat untuk mengenal pasti kehidupan kulat dalam PCR menjadikannya lebih sesuai untuk menilai keberkesanan biosida. Jadi, kajian ini bertujuan menilai keupayaan antikulat biosida, iaitu salisilat zink (ZS), kalsium benzoat (CB) dan kalium sorbate (KS) untuk merawat pencemaran kulat dalaman melalui perkembangan *limiting dilution viability-PCR* (*vPCR*). ZS, CB dan KS dipilih kerana telah dibuktikan berkesan terhadap kulat berasaskan air dalaman. Pensampelan udara dalaman mendedahkan bahawa bangunan pendidikan tinggi pengajaran komputer (Bangunan A) dan kejuruteraan awam (Bangunan B) tercemar oleh 509 CFU/m³ dan 805.7 CFU/m³ kulat udara dalaman masing-masing. Dua kulat dalaman, *Talaromyces* spp. dan *Aspergillus niger* telah dikenalpasti dan dirawat dengan biosida kerana berisiko berbahaya terhadap kesihatan manusia. Pengkulturan konvensional dan *limiting dilution vPCR* digunakan seterusnya untuk menilai kecekapan biosida. *Limiting dilution vPCR* telah diwujudkan dengan memasukkan pra-rawatan propidium monoazide (PMA) sebelum pengekstrakan asid deoksiribonukleik (DNA) dan pencairan bersiri DNA dalam PCR. Pendekatan ini terbukti berkesan untuk menilai kecekapan biosida untuk merawat kulat dalaman. KS ditunjukkan berkecekapan terbaik (100%) untuk mencegah pertumbuhan *Talaromyces* spp, diikuti oleh ZS (80.8%) dan CB (tiada kesan). KS juga menunjukkan kecekapan terbaik terhadap *A. niger* (100%) pada peringkat awal kajian tetapi kesannya menurun dengan masa. ZS menunjukkan kecekapan 66.67% terhadap kultur *A. niger* Hari-9. Keputusan yang tidak sejajar telah ditunjukkan oleh kaedah pengkulturan konvensional. Kajian ini telah menyediakan pendekatan (*limiting dilution vPCR*) yang lebih murah, tepat dan sesuai untuk menentukan kecekapan rawatan kulat dalaman berbanding kaedah pengkulturan.

TABLE OF CONTENT

TITLE	i
DECLARATION	ii
ACKNOWLEDGEMENT	iii
ABSTRACT	iv
ABSTRAK	v
TABLE OF CONTENT	vi
LIST OF TABLES	xii
LIST OF FIGURES	xv
LIST OF SYMBOLS AND ABBREVIATIONS	xx
LIST OF APPENDICES	xxiii
CHAPTER 1 INTRODUCTION	
1.1 Background of the study	1
1.2 Problem statement	4
1.3 Objectives of the investigation	5
1.4 Research questions	5
1.5 Research hypotheses	6
1.6 Scope of the investigation	7
1.7 Significance of the study	8
CHAPTER 2 LITERATURE REVIEW	
2.1 Introduction	9
2.2 Indoor environmental quality (IEQ) and indoor air quality (IAQ)	9
2.3 Standard guidelines on IAQ	11
2.3.1 Industry code of practice on indoor air quality, Malaysia	11
2.4 Sources of contamination of IAQ	13
2.4.1 Physical factors	14
2.4.2 Biological factors	15
2.5 Hazards associated with indoor fungal exposure	15

2.6	Types of indoor air contaminants	17
2.6.1	Physical risk factors	18
2.6.2	Chemical or gaseous contaminants	18
2.6.3	Particulate contaminants	19
2.6.4	Biological contaminants	19
2.7	Fungi	19
2.7.1	Properties of fungi	22
2.7.2	Growth properties of fungi	23
2.8	Isolated indoor fungi species	24
2.8.1	<i>Talaromyces</i> spp.	24
2.8.1.1	<i>Talaromyces</i> and its adverse health effect	26
2.8.2	<i>Aspergillus niger</i>	27
2.8.2.1	<i>Aspergillus</i> species and its adverse health effect	28
2.9	Malaysia climate and its effects to the buildings	29
2.9.1	Shortcomings of building that lead to microbial growth	29
2.10	Control and prevention methods for IAQ contaminations	30
2.10.1	Bioactive compounds with antifungal properties	31
2.10.1.1	Phenolics and phenolic acids	33
2.10.1.2	Potassium sorbate	34
2.10.1.3	Zinc salicylate	37
2.10.1.4	Calcium benzoate	38
2.11	Microbe monitoring and product evolution	40
2.11.1	Source sampling and air sampling	40
2.11.1.1	Culturing media and limitation of cultivation	41
2.11.2	Polymerase chain reaction (PCR)	42
2.11.2.1	Detection of microorganism	43
2.11.2.2	Detection of only live microorganism	47
2.11.2.3	Identification of species	52
2.11.2.4	PCR primer selection for <i>Talaromyces</i> spp.	52
2.11.2.5	PCR primer selection for <i>Aspergillus niger</i>	55
2.11.2.6	The use of ITS primers in fungi species identification	58
2.11.2.7	Enumeration of the species	61
2.12	Summary	61

CHAPTER 3 MATERIALS AND METHODS

3.1	Research methodology overview	63
3.2	Selection of sampling sites	63
3.3	Selection of bio-active compounds	65
3.4	Collection of sample	65
3.4.1	Preparation of media agar	66
3.4.2	Collection of air samples	66
3.4.3	Collection of surface swab samples	67
3.5	Collection of data of thermo environment	67
3.6	Measurement of microorganisms	68
3.6.1	Culture of airborne samples	68
3.6.2	Culture of surface swab samples	68
3.6.3	Enumeration of samples	69
3.7	Data analysis	69
3.8	Isolation, characterization and identification of indoor fungi	70
3.9	Laboratory-scale investigation: preparation of spore suspension	70
3.10	Preparation of biocide-treated indoor fungal species	72
3.11	PMA treatment and photo-activation of PMA	72
3.12	Isolation of total DNA from indoor fungal cell	73
3.13	Agarose gel electrophoresis (AGE)	74
3.14	Preparation of PCR primers	74
3.15	Polymerase Chain Reaction (PCR)	75
3.15.1	PCR amplification and optimization of the ITS-region of <i>Talaromyces</i> spp.	77
3.15.1.1	Optimization of PCR primer concentrations	77
3.15.1.2	Other PCR optimizations	77
3.15.2	PCR optimization of the ITS-region in rDNA gene of <i>A. niger</i>	78
3.16	Validation of PCR and vPCR	78
3.17	Quantification of PCR products using limiting dilution PCR technique	79
3.18	Spectrophotometric determination of DNA concentration	80
3.19	Antimicrobial activity test using cultivation methods	80
3.19.1	Direct biocidal test	80

3.19.2	<i>In vitro</i> inhibitory assay	81
3.20	Summary	81
CHAPTER 4 CHARACTERIZATION OF INDOOR SURFACE FUNGI		
4.1	Preliminary site inspection and data collection	84
4.2	Enumeration of surface swab samples by cultivation	86
4.3	Comparison of indoor and outdoor fungal swab samples	90
4.3.1	Concentrations of indoor and outdoor surface-growing fungi at higher educational building of computer studies	90
4.3.2	Concentrations of indoor and outdoor surface-growing fungi at higher educational engineering-based building	91
4.3.3	Total surface-growing fungi concentrations indoors and outdoors	92
4.4	Species identification of the isolated indoor wall surface fungi	94
4.4.1	Indoor wall surface fungus S1, <i>Sporisorium</i> spp.	95
4.4.2	Indoor wall surface fungus S2, <i>Talaromyces</i> spp.	96
CHAPTER 5 CHARACTERIZATION OF INDOOR MICROBIAL AIR QUALITY		
5.1	Enumeration of indoor air samples	98
5.2	Species identification of the isolated indoor airborne fungi, <i>Aspergillus niger</i>	102
CHAPTER 6 TECHNIQUE DEVELOPMENT OF LIMITING DILUTION vPCR		
6.1	Development of PCR protocol for indoor surface-growing fungi, <i>Talaromyces</i> spp.	106
6.1.1	Optimization of PCR primer concentrations	106
6.1.2	Optimization of cell wall disruption during DNA extraction	108
6.1.3	Optimization of number of PCR cycle	109
6.2	Development of PCR protocol for indoor airborne fungi, <i>Aspergillus niger</i>	111
6.2.1	PCR Optimization of the ITS-region in rDNA gene of <i>A. niger</i>	112
6.3	Validation of developed PCR protocols	114

6.4	Development of limiting dilution viability PCR	115
6.4.1	Limiting dilution PCR	116
6.4.2	Limiting dilution viability PCR	117
6.4.3	Validation of developed limiting dilution viability PCR	120

CHAPTER 7 BIOCIDES AGAINST INDOOR FUNGI BY USING LIMITING DILUTION vPCR

7.1	Introduction	122
7.2	Estimation of initial DNA concentration of ITS region of fungi cultures	123
7.2.1	Biocides-treated <i>Talaromyces</i> spp. cultures	130
7.2.2	Biocides-treated <i>Aspergillus niger</i> cultures	135
7.3	Effectiveness of each biocide against indoor fungi	140
7.3.1	Effectiveness of each biocide against <i>Talaromyces</i> spp. without viability information	140
7.3.2	Effectiveness of each biocide against <i>Talaromyces</i> spp. with viability information	142
7.3.3	Effectiveness of each biocide against <i>Aspergillus niger</i> without viability information	144
7.3.4	Effectiveness of each biocide against <i>Aspergillus niger</i> with viability information	146

CHAPTER 8 POTASSIUM SORBATE AGAINST AIRBORNE FUNGI USING CULTIVATION METHODS

8.1	Effectiveness of potassium sorbate against indoor airborne fungi using direct biocidal test	150
8.2	Effectiveness of potassium sorbate against isolated indoor airborne fungus using <i>in-vitro</i> inhibitory assay	156

CHAPTER 9 COMPARISON OF CONVENTIONAL AND PCR METHODS

9.1	Introduction	159
9.2	Comparison of growth estimation methods	159
9.3	Overall comparison of growth estimation methods for indoor fungi	162

CHAPTER 10 CONCLUSIONS AND RECOMMENDATIONS

10.1	Conclusions	165
10.2	Recommendations	170

REFERENCES

172

APPENDICES

201

VITAE

217



LIST OF TABLES

Table	Page
2.1 Acceptable range for specific parameters of IAQ (adapted from DOSH, 2010)	18
2.2 Schematic review of studies about indoor fungi found in Malaysia and Southeast Asia countries	21
2.3 <i>Talaromyces</i> -produced mycotoxins	26
2.4 Summary of bio-active compounds application for antifungal purposes	33
2.5 The usage of potassium sorbate as antifungal agents	36
2.6 The usage of zinc salicylate as antifungal agents	38
2.7 The usage of calcium benzoate as antifungal agents	39
2.8 The application of PCR in IAQ and environmental studies	45
2.9 Comparison of properties of different DNA-intercalating agents in vPCR	49
2.10 The application of PMA in IAQ and environmental studies	51
2.11 Summary review about selection of primer of <i>Talaromyces</i> spp.	53
2.12 Summary review about selection of primer of <i>Aspergillus niger</i>	56
2.13 PCR primer pairs used in IAQ studies	59
2.14 Sequence of ITS primers	61
3.1 Selected sampling sites and respective labels in this study	65
3.2 Biocides and their labels used in this study	72
3.3 Optimization step of PCR with primer ITS1/NIG	78
3.4 Summary of conducted testing	82

4.1	Range of humidity, air velocity and temperature at the sampling sites	85
4.2	Concentrations of surface-growing fungi detected at Building A	91
4.3	Concentrations of surface-growing fungi detected at Building B	91
4.4	Total surface-growing fungi concentrations indoors and outdoors	92
5.1	The IAQ physical parameters at Building A and Building B versus ICOP-IAQ (2010) standard	99
6.1	Results of optimization of PCR with primer ITS1/NIG	111
6.2	Developed PCR protocols for <i>Talaromyces</i> spp. and <i>A. niger</i>	114
6.3	Developed PCR conditions for <i>Talaromyces</i> spp. and <i>A. niger</i>	114
7.1	Estimated initial DNA concentration of ITS region of <i>Talaromyces</i> spp. Day 6 control cultures	127
7.2	Estimated initial DNA concentration of ITS region of <i>Talaromyces</i> spp. Day 9 control cultures	128
7.3	Estimated initial DNA concentration of ITS region of untreated <i>A. niger</i> Day 6 cultures	130
7.4	Estimated initial DNA concentration of ITS region of untreated <i>A. niger</i> Day 9 cultures	130
7.5	Estimated initial DNA concentration of ITS region of KS-treated <i>Talaromyces</i> spp. Day 6 cultures	131
7.6	Estimated initial DNA concentration of ITS region of KS-treated <i>Talaromyces</i> spp. Day 9 cultures	131
7.7	Estimated initial DNA concentration of ITS region of ZS-treated <i>Talaromyces</i> spp. Day 6 cultures	132
7.8	Estimated initial DNA concentration of ITS region of ZS-treated <i>Talaromyces</i> spp. Day 9 cultures	133
7.9	Estimated initial DNA concentration of ITS region of CB-treated <i>Talaromyces</i> spp. Day 6 cultures	134

7.10	Estimated initial DNA concentration of ITS region of CB-treated <i>Talaromyces</i> spp. Day 9 cultures	134
7.11	Estimated initial DNA concentration of ITS region of KS-treated <i>A. niger</i> Day 6 cultures	136
7.12	Estimated initial DNA concentration of ITS region of KS-treated <i>A. niger</i> Day 9 cultures	136
7.13	Estimated initial DNA concentration of ITS region of ZS-treated <i>A. niger</i> Day 6 cultures	137
7.14	Estimated initial DNA concentration of ITS region of ZS-treated <i>A. niger</i> Day 9 cultures	138
7.15	Estimated initial DNA concentration of ITS region of CB-treated <i>A. niger</i> Day 6 cultures	139
7.16	Estimated initial DNA concentration of ITS region of CB-treated <i>A. niger</i> Day 9 cultures	139
7.17	Effectiveness of biocides against indoor fungi	140



LIST OF FIGURES

Figure	Page
2.1 The IAQ complaints and investigation process (adapted from DOSH, 2010)	12
2.2 Examples of common naturally occurring phenolic acids	34
2.3 Structural formula of potassium sorbate	35
2.4 Structural formula of zinc salicylate	37
2.5 Structural formula of calcium benzoate	39
2.6 Detection of only live cells using PMA dye	49
2.7 Locations of ITS PCR primers on nuclear rDNAs	60
3.1 Research methodology overview	64
3.2 The Anderson single stage air impactor with high flow pump (SKC Inc, USA)	66
3.3 Collection of air samples at a sampling site	67
3.4 Environment tester equipment to measure temperature, air velocity and relative humidity (Lutron LM-8000, Taiwan)	68
3.5 The grid view and main square of a Neubauer chamber	71
3.6 Photo-activation of PMA-treated fungi cultures under 500 W halogen light	73
3.7 The Gel Doc™ XR+ Documentation System used for gel image visualisation in this study	75
3.8 The thermal cycler (C1000) used in the PCR studies	76

4.1	Mould growth at the sampling sites. (a) Male prayer room (MPR) of Building A was found to have water-leakage problems and visible moulds were observed throughout the study especially at the ceiling boards. (b) Female prayer room (FPR) of Building A was observed with serious visible moulds growth	85
4.2	Growth of indoor surface fungi sample in MPR and FPR, Building A	88
4.3	Growth of indoor surface fungi sample in MR, Building B	89
4.4	Layouts of the sampling sites (a) MPR and FPR (b) MR	90
4.5	Isolated indoor surface swab samples of fungus (a) S1 (b) S2	94
4.6	Neighbour-joining tree of the ITS sequence of indoor surface swab sample of fungus, S1 and top 10 hits Blast Results against NCBI Nucleotide Collection (nr/nt) Database by NCBI Blast Tree Method	95
4.7	Phylogenetic tree – neighbour joining analysis of the ITS sequence of the isolated indoor wall surface fungus sample, S2 to the top 10 hit Blast Results against NCBI Nucleotide Collection (nr/nt) Database by NCBI Blast Tree Method	96
5.1	Indoor air sampling for (a) fungus and (b) bacteria compared to ICOP-IAQ (2010) standard guidelines and outdoors	101
5.2	Isolated indoor airborne samples of fungus (a) FBA (b) FBB	103
5.3	Neighbour-joining tree of the ITS sequence of indoor airborne sample of fungus, FBA and FBB and top 10 hits Blast Results against NCBI Nucleotide Collection (nr/nt) Database by NCBI Blast Tree Method	103
6.1	Optimization of PCR primer concentrations in PCR of ITS region of <i>Talaromyces</i> spp.	107
6.2	1.0% AGE of PCR products of ITS region of <i>Talaromyces</i> spp. after optimization of cell wall disruption	108

6.3	Optimization of PCR cycles to 35 cycles in PCR of ITS region of <i>Talaromyces</i> spp.	109
6.4	Optimization of PCR cycles to 40 cycles in PCR of ITS region of <i>Talaromyces</i> spp.	110
6.5	Optimization of dNTPs mix concentration in PCR of ITS region of <i>A. niger</i>	112
6.6	1.0% Agarose gel electrophoresis of PCR products of ITS region of <i>A. niger</i> after optimizations	113
6.7	Validation of optimized PCR protocols (a) <i>Talaromyces</i> spp. (b) <i>A. niger</i>	115
6.8	PCR products of a 10-fold DNA dilution series of <i>Talaromyces</i> spp.	116
6.9	PCR products of a 10-fold DNA dilution series of <i>A. niger</i>	117
6.10	1.0% AGE of limiting-dilution vPCR products of ITS region of the <i>Talaromyces</i> spp. Day 3 cultures	118
6.11	1.0% AGE of limiting-dilution viability-PCR products of ITS region of the <i>A. niger</i> Day 3 cultures (PMA-untreated)	119
6.12	1.0% AGE of limiting-dilution viability-PCR products of ITS region of the <i>A. niger</i> Day 3 cultures (PMA-treated)	119
6.13	1.0% AGE of validation of limiting-dilution viability-PCR for <i>Talaromyces</i> spp.	121
6.14	1.0% AGE of validation of limiting-dilution viability-PCR for <i>A. niger</i>	121
7.1	Day 6 triplicates of <i>Talaromyces</i> spp. culture in biocides-treated MEB and control MEB (a) replicate 1 and 2 (b) replicate 3	123
7.2	Day 9 triplicates of <i>Talaromyces</i> spp. culture in biocides-treated MEB and control MEB (a) replicate 1 (b) replicate 2 (c) replicate 3	124
7.3	Growth of the <i>Talaromyces</i> spp. cultures	124
7.4	Day 6 triplicates of <i>A. niger</i> culture in biocides-treated MEB and control MEB (a) replicate 1 (b) replicate 2 (c) replicate 3	125

7.5	Day 9 triplicates of <i>A. niger</i> culture in biocides-treated MEB and control MEB (a) replicate 1 (b) replicate 2 (c) replicate 3	125
7.6	Growth of the <i>A. niger</i> cultures	125
7.7	The trend of averaged estimated initial DNA concentration of control <i>Talaromyces</i> spp. cultures	128
7.8	The trend of averaged estimated initial DNA concentration of control <i>A. niger</i> cultures	129
7.9	The trend of averaged estimated initial DNA concentration of KS-treated <i>Talaromyces</i> spp. cultures	132
7.10	The trend of averaged estimated initial DNA concentration of ZS-treated <i>Talaromyces</i> spp. cultures	133
7.11	The trend of averaged estimated initial DNA concentration of CB-treated <i>Talaromyces</i> spp. cultures	135
7.12	The trend of averaged estimated initial DNA concentration of KS-treated <i>A. niger</i> cultures	136
7.13	The trend of averaged estimated initial DNA concentration of ZS-treated <i>A. niger</i> cultures	138
7.14	The trend of averaged estimated initial DNA concentration of CB-treated <i>A. niger</i> cultures	139
7.15	Comparison of the amount of DNA of different <i>Talaromyces</i> spp. cultures without viability information on Day 6 and Day 9	141
7.16	Effectiveness of different biocides against <i>Talaromyces</i> spp. without viability information (without PMA treatment)	142
7.17	Comparison of the amount of DNA of different <i>Talaromyces</i> spp. cultures with viability information on Day 6 and Day 9	143
7.18	Effectiveness of different biocides against <i>Talaromyces</i> spp. with viability information	143
7.19	Comparison of the amount of DNA of different <i>A. niger</i> cultures without viability information on Day 6 and Day 9	145

7.20	Effectiveness of different biocides against <i>A. niger</i> without viability information	145
7.21	Comparison of the amount of DNA of different <i>A. niger</i> cultures with viability information on Day 6 and Day 9	146
7.22	Effectiveness of different biocides against <i>A. niger</i> with viability information	147
8.1	Viability of total airborne fungi at Building A (computer studies building) on KS-treated and untreated MEA	151
8.2	Viability of total airborne fungi at Building B (engineering studies building) on KS-treated and untreated MEA	152
8.3	Average viability of total airborne fungi at Building A & B on KS-treated and untreated MEA	153
8.4	The effectiveness of potassium sorbate treatment for total both samples (Building A & B)	154
8.5	Measurement of the growth diameter of the isolated fungus from Building A during biocide treatment	156
8.6	<i>In-vitro</i> inhibitory assay for sample A: Untreated fungus (a)–(b) and KS-treated fungus (c)–(d) after 7 days at 37°C incubation	157
8.7	Total measurement of the growth diameter of the isolated fungus from Building B during potassium sorbate treatment	158
9.1	Comparison of methods used to assess the effectiveness of biocide against fungi obtained in building of computer studies (Building A)	160
9.2	Comparison of methods used to assess the effectiveness of biocide against fungi obtained in building of engineering studies (Building B)	161
9.3	Overall comparison of growth estimation methods to assess the effectiveness of biocide to treat indoor fungi	162
9.4	The application of limiting dilution vPCR to treat indoor fungal contamination	164

LIST OF SYMBOLS AND ABBREVIATIONS

>	more than
%	percent
&	and
<	smaller than
×	times
µg	microgram
µl	microliter
µM	microMolar
<i>A. niger</i>	<i>Aspergillus niger</i>
AGE	agarose gel electrophoresis
AIDS	acquired immune deficiency syndrome
ANOVA	Analysis of variance
ASHRAE	American Society for Heating, Refrigerating and Air- Conditioning Engineers
b.p.	base pair
BLAST	Basic Local Alignment Search Tool
BRI	building related illnesses
C1	complained prayer room
C2	complained lecturer room 146
CB	calcium benzoate
CFU	colony forming unit
cm ²	square centimeter
dH ₂ O	sterile distilled water
DNA	deoxyribonucleic acid
dNTP	deoxynucleotide triphosphate
DOSH	Department of Occupational Safety and Health
EMA	ethidium monoazide

EPA	Environmental Protection Agency
<i>et al.</i>	and others
EtBr	ethidium bromide
FPR	female prayer room
g	gram
HVAC	heating, ventilation and air conditioning
i.e.	that is
IAQ	indoor air quality
ICOP-IAQ	industry code of practice on indoor air quality
IEQ	indoor environmental quality
ITS	internal transcribed spacer
JKPPU	Research Management Committees of UTHM
KS	potassium sorbate
L	Litre
m ³	cubic meter
Max	maximum
MEA	malt extract agar
MEB	malt extract broth
mg	milligrams
Mg ²⁺	magnesium ion
Min	minimum
min	minutes
mL	milliliter
mM	milliMolar
mm	millimeter
MPR	male prayer room
MVAC	mechanical ventilating and air conditioning
NaCl	sodium chloride
NC1	non-complained lecturer rooms' lobby
NC2	non-complained chemical engineering laboratory
NCBI	National Center for Biotechnology Information
NIOSH	National Institute of Occupational Safety and Health
NMAM	NIOSH Manual Analytical Method
NYCOSH	New York Committee for Occupational Safety and Health

°C	degree Celsius
PCR	polymerase chain reaction
PMA	propidium monoazide
rDNA	ribosomal deoxyribonucleic acid
RNA	ribonucleic acid
RNase	Ribonuclease
rpm	rotation per minute
rRNA	ribosomal ribonucleic acid
s	seconds
SBS	sick building syndrome
spp.	species
SVOCs	semi volatile organic compounds
TAE	tris-acetate-EDTA
TSA	Tryptcase Soy Agar
U	Unit
US	United States
UTHM	Universiti Tun Hussin Onn Malaysia
UV	ultraviolet
V	volt
VOCs	volatile organic compounds
vPCR	viability polymerase chain reaction
W	watt
w/v	weight per volume
ZS	zinc salicylate

LIST OF APPENDICES

A	Limiting factor of untreated <i>Talaromyces</i> spp. cultures	201
B	Limiting factor of KS-treated <i>Talaromyces</i> spp. cultures	203
C	Limiting factor of ZS-treated <i>Talaromyces</i> spp. cultures	205
D	Limiting factor of CB-treated <i>Talaromyces</i> spp. cultures	207
E	Limiting factor of untreated <i>Aspergillus niger</i> cultures	209
F	Limiting factor of KS-treated <i>Aspergillus niger</i> cultures	211
G	Limiting factor of ZS-treated <i>Aspergillus niger</i> cultures	212
H	Limiting factor of CB-treated <i>Aspergillus niger</i> cultures	214
I	Failed PCRs of ITS region of <i>A. niger</i>	216



CHAPTER 1

INTRODUCTION

1.1 Background of the study

Indoor environmental quality (IEQ) and indoor air quality (IAQ) are very important to health since most of the civilized people stay indoor, where air pollution can be far more severe than outdoor nowadays, most of the time (Orwell *et al.*, 2004). The IAQ in any building can be compromised by indoor air pollutants such as microbial contaminants (fungi, bacteria), chemicals (such as carbon monoxide, formaldehyde), allergens, or any mass or energy stressor that can induce health effects. Indoor fungal pollution can bring adverse effects to human health. It is caused by hundreds of species of a typical fungi growing indoor when sufficient moisture is available. It has been reported that dampness and moulds observations are associated with building-related health problems (Pitkaranta *et al.*, 2008). According to Pasanen, Lappalainen and Pasanen (1996), nearly 10% of people worldwide have fungal allergy. Plenty of studies pointed out that these fungi are able to produce various kinds of mycotoxins which in turn can cause adverse health effects such as asthma, headache, hypersensitive pneumonia, mucous membrane irritation, damage to body systems. However, very few studies have been done to assess indoor fungal contamination in Malaysia and even Southeast Asian countries which share common hot and humid climate (Lian, Inangda & Ramly, 2007; Ismail, Deros & Leman, 2010; Wardah *et al.*, 2011; Hussin *et al.*, 2011, Sekhar *et al.*, 2003; Luksamijarulkul *et al.*, 2009). This is a critical issue as many experts have concluded that the fungal distribution and composition is geographically patterned which make the assessment is a must at every location to ensure the safety of indoor occupants and users (Amend *et al.*,

2010). To the best of the author's knowledge, the studies of indoor fungi contamination in the higher educational building in Malaysia had not been studied before. Hence, the aim of this study is to develop a method to control the growth of specific indoor fungi species found in the higher educational buildings in Malaysia.

Researchers showed that cleaning efforts or remediation were effective in the removal of the fungal problem contributed by indoor mould growth sources (Pitkaranta *et al.*, 2011). Some studies have shown that mould exposure and related health symptoms can be reduced by removing the underlying cause of water accumulation and mould growth (Lee *et al.*, 2006). However, normal cleaning approaches cannot provide effective and sustainable solution to the fungal contamination as Pitkaranta *et al.* (2011) suggested that fungal particles aerosolized and spread during remediation. Furthermore, continuing replacing the damaged building materials is neither cost-effective nor environmental friendly. Moreover, the normal remediation approaches are not practical in Malaysia because of the high humidity contents in the weather throughout the years. Hence, there is a significant need for a sustainable, low cost and environmental friendly approach to address the IAQ problems. Researchers have sought many different methods of remediation to solve the indoor microbial contamination. Chakravarty and Korva (2013) have found that the growth of fungi was significantly inhibited by five types of antifungal agents (Sanimaster, hydrogen peroxide, isopropyl alcohol, bleach, and Sporicidin) but their effects reduced with time. This phenomenon suggests that the effects of these chemicals are not long-lasting. Moreover, treating the microbes with synthetic fungicides is harmful to the environment and human beings (Verma *et al.*, 2008).

Biocides which are rich in bioactive compounds are safer and more environment-friendly alternatives for indoor fungi control. They have long been used as antimicrobial agents especially in food and pharmaceutical industry. It has given us an insight to apply these substances to treat the indoor microbial contamination as Bellotti *et al.* (2013) has reported that some biocides from food industry, zinc salicylate, calcium benzoate and potassium sorbate, can be used to control the growth of *Chaetomium globosum* and *Alternaria alternata* in indoor waterborne coatings. However, the usage of these natural compounds in the realm of indoor environment and IAQ has not been given full address yet. The indoor fungi isolated from the contaminated higher educational buildings in this study, *Aspergillus niger* (an indoor airborne fungus) and *Talaromyces* spp. (an indoor surface-growing fungus) had not

REFERENCES

- Abarca, M. L., Bragulat, M. R., Castella, G., & Cabanes, F. J. (1994). Ochratoxin A production by strains of *Aspergillus niger* var. *niger*. *Applied and Environmental Microbiology*, 60(7), 2650-2652.
- Abarca, M. L., Accensi, F., Cano, J., & Cabañes, F. J. (2004). Taxonomy and significance of black aspergilli. *Antonie van Leeuwenhoek*, 86(1), 33-49.
- Abbott, S. P. (2002). Mycotoxins and indoor molds. *Indoor Environment Connections*, 3(4), 14-24.
- Abidin, A.S.Z., Leman, A.M., & Noraini, N.M.R. (2011) Background level of microbe in new office building. *J. Occu. Safety & Health*, 8, 25-34.
- Accensi, F., Cano, J., Figuera, L., Abarca, M. L., & Cabanes, F. J. (1999). New PCR method to differentiate species in the *Aspergillus niger* aggregate. *FEMS Microbiology Letters*, 180(2), 191–196.
- Accensi, F., Abarca, M. L., Cano, J., Figuera, L., & Cabanes, F. J. (2001). Distribution of ochratoxin A producing strains in the *A. niger* aggregate. *Antonie van Leeuwenhoek*, 79(3-4), 365-370.
- Agusti, G., Fittipaldi, M., Morat ó J., & Codony, F. (2013). Viable quantitative PCR for assessing the response of *Candida albicans* to antifungal treatment. *Applied microbiology and biotechnology*, 97(1), 341-349.

Alvarez, A.J., Buttner, M. P., Toranzos, G.A., Dvorsky E.A., Toro, A., Heikes, T.B., Mertikas, L.E., & Stetzenbach, L.D. (1994). The use of solid-phase polymerase chain reaction for the enhanced detection of airborne microorganisms. *Appl. Environ. Microbiol.* 60, 374–376.

Alvarez, A.J., Buttner, M. P., & Stetzenbach, L. D. (1995). PCR for bioaerosol monitoring: sensitivity and environmental interference. *Appl. Environ. Microbiol.* 61, 3639-3644.

Alvarez, G., González, M., Isabal, S., Blanc, V., & León, R. (2013). Method to quantify live and dead cells in multi-species oral biofilm by real-time PCR with propidium monoazide. *AMB Express*, 3(1), 1-8.

Amann, R. I., Ludwig, W. & Schleifer, K. H. (1995). Phylogenetic identification and in situ detection of individual microbial cells without cultivation. *Microbiol. Rev.* 59, 143–169.

Ambu, S., Chu, W. L., Mak, J. W., Wong, S. F., Chan, L. L., & Wong, S. T. (2008). Environmental health and building related illnesses. *Journal of Science, Medicine and Education*, 2(1), 11-18.

Amend, A.S., Seifert, K.A., Samson, R., & Bruns, T.D. (2010). Indoor fungal composition is geographically patterned and more diverse in temperate zones than in the tropics. *PNAS* 107(310), 13748–13753. www.pnas.org/cgi/doi/10.1073/pnas.1000454107

Anderson, G., & Palombo, E. A. (2009). Microbial contamination of computer keyboards in a university setting. *American journal of infection control*, 37(6), 507-509.

Andorrà I, Esteve-Zarzoso B, Guillamón JM, & Mas, A. (2010) Determination of viable wine yeast using DNA binding dyes and quantitative PCR. *Int J Food Microbiol* 144, 257 – 262.

- Anjaneyulu, Y., & Raman, N. B. (1985). Physicochemical and antimicrobial activity studies on the ternary complexes of zinc (II) with 8-hydroxyquinoline and salicylic acids. *Indian J. Microbiol*, 25(3-4), 125-128.
- Aneja, K. R. (2003). *Experiments in microbiology, plant pathology and biotechnology*. New Age International.
- ASHRAE. 2007. *Ventilation for Acceptable Indoor Air Quality Standard 62.1*. United State. associated with some toxic fungi and their mycotoxins. *Analyst*, 121, 1949-1953.
- Balasundram, N., Sundram, K., & Samman, S. (2006) Phenolic compounds in plants and agriindustrial by-products: antioxidant activity, occurrence, and potential uses. *Food Chem*, 99, 191–203.
- Balouiri, M., Sadiki, M., & Ibensouda, S. K. (2016). Methods for in vitro evaluating antimicrobial activity: A review. *Journal of Pharmaceutical Analysis*, 6(2), 71-79.
- Barbas, C. F., Burton, D. R., Scott, J. K., & Silverman, G. J. (2007). Quantitation of DNA and RNA. *Cold Spring Harbor Protocols*, 2007(11), pdb-ip47
- Baughman, A., & Arens, E. A. (1996). Indoor humidity and human health--Part I: Literature review of health effects of humidity-influenced indoor pollutants. *ASHRAE Transactions*, 102.
- Bej, A. K., Ng, W.Y., Morgan, S., Jones, D., & Mahbubani, M. H. (1996). Detection of viable *Vibrio cholerae* by reverse-transcriptase polymerase chain reaction (RT-PCR). *Mol. Biotechnol.* 5, 1–10.
- Benjamin, C. R. (1955). Ascocarps of *Aspergillus* and *Penicillium*. *Mycologia*, 47(5), 669-687.

- Bellanger, A. P., Reboux, G., Roussel, S., Grenouillet, F., Didier-Scherer, E., Dalphin, J. C., & Millon, L. (2009). Indoor fungal contamination of moisture-damaged and allergic patient housing analysed using real-time PCR. *Letters in applied microbiology*, 49(2), 260-266.
- Bellotti, N., Salvatore, L., Deya, C., Del Panno, M.T., del Amo, B., & Rogmanli, R. (2013). The application of bioactive compounds from the food industry to control mold growth in indoor waterborne coatings. *Colloids and Surfaces B: Biointerfaces*, 104, 140-114.
- Berardi, B.M., Leoni, E., Marchesini, B., Cascella, D., & Raffil, G.B. (1991). Indoor climate and air quality in new offices: effects of a reduced air-exchange rate. *International Archives of Occupational and Environmental Health*, 63, 233–239.
- Berbee, M. L., & Taylor, J. W. (1993). Ascomycete relationships: dating the origin of asexual lineages with 18S ribosomal RNA gene sequence data. *The Fungal Holomorph: mitotic, meiotic and pleomorphic speciation in fungal systematics*, 67-78.
- Bleve, G., Rizzotti, L., Dellaglio, F., & Torriani, S., (2003). Development of reverse transcription (RT)-PCR and real-time RT-PCR assays for rapid detection and quantification of viable yeasts and molds contaminating yogurts and pasteurized food products. *Appl. Environ. Microbiol.* 69, 4116–4122.
- Bodey, G., Bueltmann, B., Duguid, W., Gibbs, D., Hanak, H., Hotchi, M., Mall, G., Martino, P., Meunier, F., Milliken, S. & Naoe, S. (1992). Fungal infections in cancer patients: an international autopsy survey. *European Journal of Clinical Microbiology and Infectious Diseases*, 11(2), 99-109.
- Borrego, S., Lavin, P., Perdomo, I., Gómez de Saravia, S., & Guiamet, P. (2012). Determination of indoor air quality in archives and biodeterioration of the documentary heritage. *ISRN microbiology*, 2012.

- Boswell, G. P., Jacobs, H., Davidson, F. A., Gadd, G. M., & Ritz, K. (2003). Growth and function of fungal mycelia in heterogeneous environments. *Bulletin of mathematical biology*, 65(3), 447.
- Brasel, T. L., D. R. Douglas, S. C. Wilson, & D. C. Straus. (2005). Detection of airborne *Stachybotrys chartarum* macrocyclic trichothecene mycotoxins on particulates smaller than conidia. *Appl. Environ. Microbiol.* 71, 114–122.
- Brescia, C. C., Griffin, S. M., Ware, M. W., Varughese, E. A., Egorov, A. I., & Villegas, E. N. (2009). *Cryptosporidium parvum* monoazide-PCR, a molecular biology-based technique for genotyping of viable *Cryptosporidium* oocysts. *Applied and environmental microbiology*, 75(21), 6856-6863.
- Bruns, T. D., White, T. J., & Taylor, J. W. (1991). Fungal molecular systematics. *Annual Review of Ecology and systematics*, 525-564.
- Bujdošová, Z., Györyová, K., Hudecová, D., Kovářová, J., & Halás, L. (2010). Preparation, spectral, thermal, and biological properties of zinc (II) 4-chloro- and 5-chlorosalicylate complexes with methyl 3-pyridylcarbamate and phenazone. *Chemical Papers*, 64(5), 584-591.
- Burge H.A., Pierson D.L., Groves T.O., Strawn K.F., & Mishra S.K. (2000). Dynamics of airborne fungal populations in a large office building. *Curr Microbiol* 40(1), 10-6.
- Buttner, M. P., Cruz-Perez, P., & Stetzenbach, L. D. (2001). Enhanced detection of surface-associated bacteria in indoor environments by quantitative PCR. *Applied and Environmental Microbiology*, 67(6), 2564-2570.
- Cai, G.H., Hashim, J.H., Hashim, Z., Ali, F., Bloom, E., Larsson, L., Lampa, E., Norback, D. (2011) Fungal DNA, allergens, mycotoxins and associations with asthmatic symptoms among pupils in schools from Johor Bahru, Malaysia. *Pediatric Allergy and Immunology*. 22, 290–297.

- Carocho, M., Barreiro, M. F., Morales, P., & Ferreira, I. C. (2014). Adding molecules to food, pros and cons: A review on synthetic and natural food additives. *Comprehensive Reviews in Food Science and Food Safety*, 13(4), 377-399.
- Castillo, F., Hernandez, D., Gallegos, G., Rodríguez, R., & Aguilar, C. N. (2012). Antifungal properties of bioactive compounds from plants. *Fungicides for plant and animal diseases. Croatia, Europe: InTech*.
- Chakraborty, B. N., Chakraborty, U., Sunar, K., & Dey, P. L. (2011). RAPD profile and rDNA sequence analysis of *Talaromyces flavus* and *Trichoderma* species. *Indian Journal of Biotechnology*, 10, 487-495.
- Chakravarty, P. & Kovar, B (2013). Evaluation of Five Antifungal Agents Used in Remediation Practices Against Six Common Indoor Fungal Species, *Journal of Occupational and Environmental Hygiene*, 10(1), D11-D16
- Chan, D. W., & Miao, H. E. (2003). A Case Study of Control Strategies and Contamination Routes of Indoor Moulds for the Hotel of Philippines. *Acta Scientiarum Naturalium Universitatis Sunyatseni*, S1.
- Chang, C. W., & Lin, M. H. (2017). Optimization of PMA-qPCR for *Staphylococcus aureus* and determination of viable bacteria in indoor air. *Indoor Air*. 28(1), 64-72.
- Chao, H.J., Schwartz, J., Milton, D.K., & Burge, H.A. (2003) The work environment and workers' health in four large office buildings. *Environ Health Perspect* 111(9), 1242-8.
- Chen, A. J., Sun, B. D., Houbraken, J., Frisvad, J. C., Yilmaz, N., Zhou, Y. G., & Samson, R. A. (2016). New *Talaromyces* species from indoor environments in China. *Studies in Mycology*, 84, 119-144.

- Christaki, E., Bonos, E., Giannenas, I., & Florou-Paneri, P. (2012). Aromatic Plants as a Source of Bioactive Compounds. *Agriculture*, 2, 228-243.
- Ciferri, O. (1999). Microbial degradation of paintings. *Appl. Environ. Microbiol.*, 65(3), 879-885.
- Clayton, E. E., Smith, T. E., Shaw, K. J., Gaines, J. G., Graham, T. W., & Yeager Jr, C.C. (1943). Fungicidal tests on blue mold (*Peronospora tabacina*) of tobacco. *Jour. Agr. Res*, 66, 261-276.
- Cojocar, G. A., & Antocea, A. O. (2012). Chemical and Biochemical Mechanisms of Preservatives Used in Wine: A Review. *dio*, 1, 100.
- Conforti, F., Menichini, F., Formisano, C., Rigano, D., Senatore, F., & Arnold, N.A. (2009). Comparative chemical composition, free radical-scavenging and cytotoxic properties of essential oils of six *Stachys* species from different regions of the Mediterranean Area. *Food Chem*, 116, 898–905.
- Cookingham, C. E., & Solomon, W. R. (1995). 10 Bioaerosol-Induced Hypersensitivity Diseases. *Bioaerosols*, 2, 205-234.
- Coombs, K. C. (2015). *The Indoor Environment of Green versus Non-Green Buildings*. (Doctoral dissertation, University of Cincinnati).
- Delgado-Viscogliosi, P., Solignac, L., & Delattre, J. M. (2009). Viability PCR, a culture-independent method for rapid and selective quantification of viable *Legionella pneumophila* cells in environmental water samples. *Applied and environmental microbiology*, 75(11), 3502-3512.
- Denning, D. W. (1996). Therapeutic outcome in invasive aspergillosis. *Clinical Infectious Diseases*, 23(3), 608-615.
- Denning, D. W. (1998). Invasive aspergillosis. *Clinical infectious diseases*, 781-803.

- DOSH (2010). *Industry Code of Practice on Indoor Air Quality 2010*, Ministry of Human Resources. Department of Occupational Safety and Health, Malaysia
- Ducel, G., Fabry, J., Nicolle, L., & World Health Organization. (2002). Prevention of hospital-acquired infections: a practical guide.
- Duru, M. E., Cakir, A., Kordali, S., Zengin, H., Harmandar, M., Izumi, S., & Hirata, T. (2003). Chemical composition and antifungal properties of essential oils of three *Pistacia* species. *Fitoterapia*, 74(1), 170-176.
- Er, C. M., Sunar, N. M., Leman, A. M., Norzila, O., Emparan, Q., Kalthsom, U., Gani, P. & Jamal, N. A. (2014). Evaluation of potassium sorbate as a biocide to reduce viability of total airborne fungi in a higher educational building of computer studies. *Mater. Sci*, 2(1), 15-19.
- Er, C. M., Sunar, N. M., Leman, A. M., & Othman, N. (2015). Direct growth inhibition assay of total airborne fungi with application of biocide-treated malt extract agar. *MethodsX*, 2, 340-344.
- Fadda, A., Barberis, A., D'Aquino, S., Palma, A., Angioni, A., Lai, F., & Schirra, M. (2015). Residue levels and performance of potassium sorbate and thiabendazole and their co-application against blue mold of apples when applied as water dip treatments at 20 or 53 ° C. *Postharvest Biology and Technology*, 106, 33-43.
- Fang, Z., Ouyang, Z., Hu, L., Wang, X., Zheng, H., & Lin, X. (2005). Culturable airborne fungi in outdoor environments in Beijing, China. *Science of the Total Environment*, 350(1-3), 47-58.
- Favero, M. S., Puleo, J. R., Marshall, J. H. & Gordon S. O. (1966). Comparative levels and types of microbial contamination detected in industrial clean rooms. *Applied microbiology*, 14(4), 539-551.

- Fischer, G., & Dott, W. (2003). Relevance of airborne fungi and their secondary metabolites for environmental, occupational and indoor hygiene. *Archives of Microbiology*, 179(2), 75-82.
- Fittipaldi, M., Rodriguez, N. J. P., Codony, F., Adrados, B., Peñuela, G. A., & Morat ó J. (2010). Discrimination of infectious bacteriophage T4 virus by propidium monoazide real-time PCR. *Journal of virological methods*, 168(1), 228-232.
- Fittipaldi, M., Nocker, A., & Codony, F. (2012). Progress in understanding preferential detection of live cells using viability dyes in combination with DNA amplification. *Journal of microbiological methods*, 91(2), 276-289.
- Fredricks, D. N., Smith, C., & Meier, A. (2005). Comparison of six DNA extraction methods for recovery of fungal DNA as assessed by quantitative PCR. *Journal of clinical microbiology*, 43(10), 5122-5128.
- Frisvad, J. C., Filtenborg, O., Samson, R. A., & Stolk, A. C. (1990). Chemotaxonomy of the genus *Talaromyces*. *Antonie van Leeuwenhoek*, 57(3), 179-189.
- Garrett, M. H., Rayment, P. R., Hooper, M. A., Abramson, M. J., & Hooper, B. M. (1998). Indoor airborne fungal spores, house dampness and associations with environmental factors and respiratory health in children. *Clinical and Experimental Allergy*, 28(4), 459-467.
- Gaskell, G. J., D. A. Carter, W. J. Britton, E. R. Tovey, F. H. Benyon, and U. Lovborg. (1997). Analysis of the internal transcribed spacer regions of ribosomal DNA in common airborne allergenic fungi. *Electrophoresis* 18, 1567–1569.
- Gaunt, I. F., Butterworth, K. R., Hardy, J., & Gangolli, S. D. (1975). Long-term toxicity of sorbic acid in the rat. *Food and cosmetics toxicology*, 13(1), 31-45.

- Goh, J. C., Juliana, J., Malina, O., Ngah, Z. U., & Norhafizalena, O. (2007). Prevalence of *Penicillium* specific Ig E level and allergy symptoms among office workers in a selected company in Bangi, Malaysia. *Trop Biomed*, 24(1), 37-46.
- Gonzalez-Salgado, A., Patiño, B., Vázquez, C., & Gonzalez-Jaén, M. T. (2005). Discrimination of *Aspergillus niger* and other *Aspergillus* species belonging to section *Nigri* by PCR assays. *FEMS microbiology letters*, 245(2), 353-361.
- Graves, D.E., Watkins, C.L., & Yielding, L.W. (1981). Ethidium bromide and its photoreactive analogues: spectroscopic analysis of deoxyribonucleic acid binding properties. *Biochemistry* 20, 1887–1892.
- Graybill, J. R. (1992). Future directions of antifungal chemotherapy. *Clinical infectious diseases*, 14(Supplement 1), S170-S181.
- Green, C.F., Scarpino, P.V., & Gibbs, S.G. (2003) Assessment and modeling of indoor fungal and bacterial concentrations. *Aerobiologia*, 19, 159-169.
- Grisoli, P., Rodolfi, M., Chiara, T., Zonta, L. A., & Dacarro, C. (2012). Evaluation of microbiological air quality and of microclimate in university classrooms. *Environmental monitoring and assessment*, 184(7), 4171-4180.
- Guynot, M. E., Ramos, A. J., Sanchis, V., & Marin, S. (2005). Study of benzoate, propionate, and sorbate salts as mould spoilage inhibitors on intermediate moisture bakery products of low pH (4.5–5.5). *International journal of food microbiology*, 101(2), 161-168
- Haleem Khan, A. A., & Mohan Karuppayil, S. (2012). Fungal pollution of indoor environments and its management. *Saudi journal of biological sciences*, 19(4), 405-426.

- Ham, S.S., Kim, S.H., Moon, S.Y., Chung, M.J., Cui, C.B., & Han, E.K. (2009) Antimutagenic effects of subfractions of Chaga mushroom (*Inonotus obliquus*) extract. *Mutat Res Gen Toxicol En*, 672, 55–59.
- Hammes, F., Berney, M., & Egli, T. (2011). Cultivation-independent assessment of bacterial viability. *Adv. Biochem. Eng. Biotechnol.* 124, 123–150.
- Handbook, D. P. (2012). Sample and Assay Technologies.
- Han, J. (2003). Antimicrobial food packaging. *Novel Food Packaging Techniques*, (i). Retrieved from http://books.google.com/books?hl=en&lr=&id=6RWkAgAAQBAJ&oi=fnd&pg=PA50&dq=Antimicrobial+food+packaging&ots=NNFYR1MFFb&sig=6Sx4_xEoxydRT2Tx5IA7dnqSgjs
- Hardin, B.D., Kelman, B.J. & Saxon, A. (2003). Adverse human health effects associated with molds in the indoor environment. *J Occup Environ Med*, 45, 470–478.
- Haugland, R.A., & Heckman J.L. (1998). Identification of putative sequence specific PCR primers for detection of the toxigenic fungal species *Stachybotrys chartarum*. *Mol Cell Probes*, 12, 387.
- Haugland, R.A., Varma, M., Wymer, L.J. & Vesper, S.J. (2004) Quantitative PCR analysis of selected *Aspergillus*, *Penicillium* and *Paecilomyces* species. *Syst Appl Microbiol*, 27, 198–210.
- Heneghan, M. N., McLoughlin, L., Murray, P. G., & Tuohy, M. G. (2007). Cloning, characterisation and expression analysis of α -glucuronidase from the thermophilic fungus *Talaromyces emersonii*. *Enzyme and Microbial Technology*, 41(6), 677-682.

- Henry, T., Iwen, P. C., & Hinrichs, S. H. (2000). Identification of *Aspergillus* species using internal transcribed spacer regions 1 and 2. *Journal of Clinical Microbiology*, 38(4), 1510-1515.
- Herrera, M. L., Vallor, A. C., Gelfond, J. A., Patterson, T. F., & Wickes, B. L. (2009). Strain-dependent variation in 18S ribosomal DNA copy numbers in *Aspergillus fumigatus*. *Journal of clinical microbiology*, 47(5), 1325-1332.
- Heydaryinia, A., Veissi, M., & Sadadi, A. (2011). A comparative study of the effects of the two preservatives, sodium benzoate and potassium sorbate on *Aspergillus niger* and *Penicillium notatum*. *Jundishapur journal of microbiology*, 4(4), 301-307.
- Hilmarsdottir, I., Meynard, J. L., Rogeaux, O., Guernonprez, G., Datry, A., Katlama, C., Brückner, G., Coutellier, A., Danis, M. & Gentilini, M. (1993). Disseminated *Penicillium marneffei* infection associated with human immunodeficiency virus: a report of two cases and a review of 35 published cases. *JAIDS Journal of Acquired Immune Deficiency Syndromes*, 6(5), 466-471.
- Hixon, S. C., White, W. E., & Yielding, K. L. (1975). Selective covalent binding of an ethidium analog to mitochondrial DNA with production of petite mutants in yeast by photoaffinity labeling. *Journal of Molecular Biology*, 92, 319-329.
- Husman, T. (1996). Health effects of indoor-air microorganisms. *Scandinavian journal of work, environment & health*, 5-13.
- Hussin, N.H.M., Sann, L.M., Shamsudin, M.N., & Hashim, Z. (2011). Characterization of Bacteria and Fungi Bioaerosol in the Indoor Air of selected Primary Schools in Malaysia. *Indoor and Built Environment*, 20(6), 607-617.
- Infection Control and Hospital Epidemiology (2006) January 2006 Issue.

- Ismail, A.R., Jusoh, N., Zulkifli, R., Sopian, K., Deros, B.M. (2009). Thermal Comfort Assessment. A Case Study at Malaysian Automotive Industry. *American Journal of Applied Sciences*, 6(8), 1495–1501.
- Ismail, S.H., Deros, B.M. & Leman, A.M. (2010). Indoor Air Quality Issues for Non-Industrial Workplace. *Int. J. Res. Rev. Appl. Sci*, 5, 235-244.
- Jayanetra, P., Nitiyanant, P., Ajello, L., Padhye, A. A., Lolekha, S., Atichartakarn, V., Vathesatogit, P., Sathaphatayavongs, B., & Prajaktam, R. (1984). Penicilliosis marneffeii in Thailand: report of five human cases. *The American journal of tropical medicine and hygiene*, 33(4), 637-644.
- Jiménez, J.P., Serrano, J., Tabernero, M., Arranz, S., Díaz-Rubio, M.E., & García-Diz, L. (2008). Effects of grape antioxidant dietary fiber in cardiovascular disease risk factors. *Nutrition*, 24, 646–53.
- John, D.S., Jonathan, M.S., & John, F.M. (2001). *Indoor Air Quality Handbook*. McGraw-Hill, United States of America.
- Jones, A. P. (1999). Indoor air quality and health. *Atmospheric environment*, 33(28), 4535-4564.
- Junhui, Z., Ruifu, Y., Fengxiang, C., Meiling, C., & Hong, C. (1997). Polymerase chain reaction analysis of laboratory generated bioaerosols. *Aerobiologia*, 13(1), 7–10.
- Karbowska-Berent, J., Górny, R. L., Strzelczyk, A. B., & Wlazło, A. (2011) Airborne and dust borne microorganisms in selected polish libraries and archives. *Building and Environment*, 46(10), 1872-1879.
- Kim, G.N., Shin, J.G., & Jang, H.D. (2009) Antioxidant and antidiabetic activity of Dangyuja (*Citrus grandis* Osbeck) extract treated with *Aspergillus saitoi*. *Food Chem.* 117, 35–41.

- Kim, S. H., Ahn, G. R., Son, S. Y., Bae, G. N., & Yun, Y. H. (2014). Mold occurring on the air cleaner high-efficiency particulate air filters used in the houses of child patients with atopic dermatitis. *Mycobiology*, 42(3), 286-290.
- Koomen, G.J., Den Blaauwen, T., Hellingwerf, K.J., Ungaro, R., & Mobashery, S. (2002). Fighting microbial resistance through development of new antimicrobial agents, directed against new specific targets. *IUPAC Project* 2002-030-1-300.
- Kozak, J. P., Gallup, J., Cummins, L. H., & Gillman, S. A. (1980). Currently available methods for home mold surveys. II. Examples of problem homes surveyed. *Annals of Allergy*, 45(3), 167-176.
- Kris-Etherton, P.M., Hecker, K.D., Bonanome, A., Coval, S.M., Binkoski, A.E., & Hilpert, K.F. (2002) Bioactive compounds in foods: their role in the prevention of cardiovascular disease and cancer. *Am J Med*, 113, 71S–88S
- Krogh, P., Hald, B., Pleština, R., & Čević, S. (1977). Balkan (endemic) nephropathy and foodborn ochratoxin A: preliminary results of a survey of foodstuffs. *Acta Pathologica Microbiologica Scandinavica Section B Microbiology*, 85(3), 238-240.
- Kuhn, D.M. & Ghannoum, M.A. (2003) Indoor mold, toxigenic fungi, and *Stachybotrys chartarum*: infectious disease perspective. *Clin Microbiol Rev*, 16, 144–172.
- Lawton, M.D., Dales, R.E., & White, J. (1998). The influence of house characteristics in a Canadian community on microbiological contamination. *Indoor Air*, 8, 2-11.
- Lee, J. L., & Levin, R. E. (2006). Use of ethidium bromide monoazide for quantification of viable and dead mixed bacterial flora from fish fillets by polymerase chain reaction. *Journal of microbiological methods*, 67(3), 456-462.

- Lee, S., Adhikari, A., Grinshpun, S., McKay, R., Shukla, R., & Reponen, T. (2006). Personal Exposure to Airborne Dust and Microorganisms in Agricultural Environments. *Journal of Occupational and Environmental Hygiene*, 3, 118-130.
- Li, C. S. (1999). Sampling performance of impactors for bacterial bioaerosols. *Aerosol Science & Technology*, 30(3), 280-287.
- Lian, K.S., Inangda, P.S.N., & Ramly, A. (2007). Sources of Indoor Air Quality Problem in a New Hospital in Malaysia. *The 6th International Conference on Indoor Air Quality, Ventilation & Energy Conservation in Buildings*, Sendai, Japan.
- Lobuglio, A. K. F., Pitt, J. I., Taylor, J. W., & Url, S. (1993). Phylogenetic Analysis of Two Ribosomal DNA Regions Indicates Multiple Independent Losses of a Sexual *Talaromyces* State among Asexual *Penicillium* Species in Subgenus *Biverticillium*. *Mycological Society of America*, 85(4), 592-604.
- LoBuglio, K. F., & Taylor, J. W. (1995). Phylogeny and PCR identification of the human pathogenic fungus *Penicillium marneffeii*. *Journal of clinical microbiology*, 33(1), 85-89.
- Loison, M. (2010). *The plant protection corner, in: A growing industry, various business models*. New Ag Internacional magazine, (February 2010). www.newaginternational.com/current/plantprotection201011.pdf
- Lou, X., Fang, Z., & Si, G. (2012). Assessment of culturable airborne bacteria in a university campus in Hangzhou, Southeast of China. *African journal of microbiology research*, 6(3), 665-673.
- Luksamijarulkul, P., Panya, N., Sujirarat, D., & Thaweboon, S. (2009). Microbial air quality and standard precaution practice in a hospital dental clinic. *Journal of the Medical Association of Thailand= Chotmaihet thangphaet*, 92, S148-155.

- Lymperopoulou, D. S., Adams, R. I., & Lindow, S. E. (2016). Contribution of vegetation to the microbial composition of nearby outdoor air. *Applied and environmental microbiology*, 82(13), 3822-3833.
- Macher, J. (Ed.) (1999). *Bioaerosols: Assessment and Control*. American Conference Governmental Industrial Hygienists ACGIH. Cincinnati.
- MacNeil, L., Kauri, T., & Robertson, W. (1995). Molecular techniques and their potential application in monitoring the microbiological quality of indoor air. *Canadian journal of microbiology*, 41(8), 657-665.
- Mahnert, A., Moissl-Eichinger, C., & Berg, G. (2015). Microbiome interplay: plants alter microbial abundance and diversity within the built environment. *Frontiers in microbiology*, 6.
- Malta-Vacas, J., Viegas, S., Sabino, R., & Viegas, C. (2012). Fungal and microbial volatile organic compounds exposure assessment in a waste sorting plant. *Journal of Toxicology and Environmental Health, Part A*, 75(22-23), 1410-1417.
- Martin, E., Kämpfer, P., & Jäckel, U. (2009). Quantification and identification of culturable airborne bacteria from duck houses. *Annals of occupational hygiene*, 54(2), 217-227.
- Martinez, C., Roux, C., Jauneau, A., & Dargent, R. (2002). The biological cycle of *Sporisorium reilianum* f. sp. *zeae*: an overview using microscopy. *Mycologia*, 94(3), 505-514.
- Martinez, R. (2011). *How to Count With a Neubauer Chamber*. Retrieved November 19, 2013, from <http://www.biocenter.helsinki.fi/bi/gbu/orf/Counting%20cells.pdf>

- Martins, S., Mussatto, S.I., Martínez-Avila, G., Montañez-Saenz, J., Aguilar, C.N., & Teixeira, J.A. (2011) Bioactive phenolic compounds: Production and extraction by solid-state fermentation. A review. *Biotechnology Advances*, 29, 365–373
- Martyny, J. (1999). Source sampling. In: Macher, J., editor. *Bioaerosols: Assessment and Control*. Cincinnati, Ohio: ACGIH, p. 12–1p. 12–5
- Marx, G., Zhou, H., Graves, D. E., & Osheroff, N. (1997). Covalent attachment of ethidium to DNA results in enhanced topoisomerase II-mediated DNA cleavage. *Biochemistry*, 36(50), 15884-15891.
- Massey, D. D., & Taneja, A. (2011). Emission and formation of fine particles from hardcopy devices: the cause of indoor air pollution. *Monit Control and Eff Air Pollut*, 9, 121-134.
- Mehyar, G. F., Al-Qadiri, H. M., Abu-Blan, H. A., & Swanson, B. G. (2011). Antifungal effectiveness of potassium sorbate incorporated in edible coatings against spoilage molds of apples, cucumbers, and tomatoes during refrigerated storage. *Journal of food science*, 76(3), M210-M217.
- Mehyar, G. F., Al-Qadiri, H. M., & Swanson, B. G. (2014). Edible coatings and retention of potassium sorbate on apples, tomatoes and cucumbers to improve antifungal activity during refrigerated storage. *Journal of food processing and preservation*, 38(1), 175-182.
- Meklin, T., Reponen, T., McKinstry, C., Cho, S. H., Grinshpun, S. A., Nevalainen, A., Vepsäläinen, A., Haugland, R.A., LeMasters, G., & Vesper, S. J. (2007). Comparison of mold concentrations quantified by MSQPCR in indoor and outdoor air sampled simultaneously. *Science of the Total Environment*, 382(1), 130-134.

Melchers, W. J., Verweij, P. E., Van Den Hurk, P., Van Belkum, A., De Pauw, B. E., Hoogkamp-Korstanje, J. A., & Meis, J. F. (1994). General primer-mediated PCR for detection of *Aspergillus* species. *Journal of Clinical Microbiology*, 32(7), 1710-1717.

Meletiadis, J., Meis, J. F., Mouton, J. W., & Verweij, P. E. (2001). Analysis of growth characteristics of filamentous fungi in different nutrient media. *Journal of Clinical Microbiology*, 39(2), 478-484.

Merck. (2005). *Microbiology manual* (12th ed.). Darmstadt: Merck KGaA, p. 344.

Miller, J.D. (1992) Fungi as contaminants in indoor air. *Atmospheric Environment. Part A. General Topics*, 26(12), 2163-2172.

Montealegre, F., Fernández, B., Delgado, A., Fernández, L., Román, A., Chardón, D. & Bayona, M. (2004). Exposure levels of asthmatic children to allergens, endotoxins, and serine proteases in a tropical environment. *Journal of Asthma*, 41(4), 485-496.

Morin, N.J., Gong, Z., & Li, X.F., (2004). Reverse transcription-multiplex PCR assay for simultaneous detection of *Escherichia coli* O157: H7, *Vibrio cholerae* O1, and *Salmonella typhi*. *Clin. Chem.* 50, 2037–2044.

Moss, M. O., & Hill, I. W. (1970). Strain variation in the production of rubratoxins by *Penicillium rubrum* Stoll. *Mycopathologia et mycologia applicata*, 40(2), 81-88.

Mukoda, T. J., Todd, L.A., & Sobsey, M.D. (1994). PCR and gene probes for detecting bioaerosols. *J. Aerosol Sci.* 25, 1523-1532.

Nayak, B. K., & Hazra, A. (2011). How to choose the right statistical test? *Indian journal of ophthalmology*, 59(2), 85.

- Nielsen, K.F. (2003). Mycotoxin production by indoor moulds. *Fungal genetics and biology*, 39(2), 103-117.
- Niemeier, R.T., Sivasubramani, S. K., Reponen, T. & Grinshpun, S. A. (2006) Assessment of Fungal Contamination in Moldy Homes: Comparison of Different Methods. *J Occup Environ Hyg*. 2006 May, 3(5), 262–273.
- Nkuipou-Kenfack, E., Engel, H., Fakih, S., & Nocker, A. (2013). Improving efficiency of viability-PCR for selective detection of live cells. *Journal of microbiological methods*, 93(1), 20-24.
- Nocker, A., & Camper, A. K. (2006). Selective removal of DNA from dead cells of mixed bacterial communities by use of ethidium monoazide. *Applied and Environmental Microbiology*, 72, 1997-2004.
- Nocker, A. & Camper, A.K. (2009). Novel approaches toward preferential detection of viable cells using nucleic acid amplification techniques. *FEMS microbiology letters*, 291(2), 137-142.
- Nocker, A., Cheung, C. Y., & Camper, A. K. (2006). Comparison of propidium monoazide with ethidium monoazide for differentiation of live vs. dead bacteria by selective removal of DNA from dead cells. *Journal of Microbiological Methods*, 67, 310-320.
- Nogva, H. K., Drømtorp, S. M., Nissen, H., & Rudi, K. (2003). Ethidium monoazide for DNA-based differentiation of viable and dead bacteria by 5'-nuclease PCR. *BioTechniques* 34, 804-813.
- Norbäck, D., Hashim, J. H., Hashim, Z., Cai, G. H., Sooria, V., Ismail, S. A., & Wieslander, G. (2017). Respiratory symptoms and fractional exhaled nitric oxide (FeNO) among students in Penang, Malaysia in relation to signs of dampness at school and fungal DNA in school dust. *Science of The Total Environment*, 577, 148-154.

NYCOSH, 2013. Methods for Evaluation of Indoor Mold Growth.

<http://nycosh.org/index.php?page=Biological> Retrieved: 01-18-2013.

Odell, E., & Pertl, C. (1995). Zinc as a growth factor for *Aspergillus* sp. and the antifungal effects of root canal sealants. *Oral Surgery, Oral Medicine, Oral Pathology, Oral Radiology, and Endodontology*, 79(1), 82-87.

Orhan, D.D., O'zcelik, B., o'zgen, S., & Ergun, F. (2010) Antibacterial, antifungal, and antiviral activities of some flavonoids. *Microbiological Research*, 165, 496—504.

Orwell, R. L., Wood, R. L., Tarran, J., Torpy, F., & Burchett, M. D. (2004) Removal of benzene by the indoor plant/substrate microcosm and implication for air quality. *Water, Air, and Soil Pollution*, 157(1-4), 193–207,

Ottow, J. C., & Glathe, H. (1968). Rose bengal-malt extract-agar, a simple medium for the simultaneous isolation and enumeration of fungi and actinomycetes from soil. *Applied microbiology*, 16(1), 170.

Ough, C. S., & Amerine, M. A. (1988). *Methods for analysis of musts and wines*. J. Wiley.

Pan, Y., & Breidt, F. (2007). Enumeration of viable *Listeria monocytogenes* cells by real-time PCR with propidium monoazide and ethidium monoazide in the presence of dead cells. *Applied and Environmental Microbiology*, 73(24), 8028-8031.

Parjo, U. K., Sunar, N. M., Leman, A. M., Gani, P., Embong, Z., & Tajudin, S. A. A. (2016). The *Aspergillus niger* growth on the treated concrete substrate using variable antifungals. *IOP Conference Series: Materials Science and Engineering*, 160(1), 012051. IOP Publishing

- Park, S. I., Stan, S. D., Daeschel, M. A., & Zhao, Y. (2005). Antifungal coatings on fresh strawberries (*Fragaria* × *ananassa*) to control mold growth during cold storage. *Journal of Food Science*, 70(4), M202-M207.
- Parvathy, K.S., Negi, P.S., & Srinivas, P. (2009) Antioxidant, antimutagenic and antibacterial activities of curcumin- β -diglucoside. *Food Chem*, 115, 265–271.
- Pasanen, A. L., Lappalainen, S., & Pasanen, P. (1996). Volatile organic metabolites associated with some toxic fungi and their mycotoxins. *Analyst*, 121(12), 1949-1953.
- Patel, B.K.R., Banerjee, D.K., & Butcher, P.D. (1993). Determination of *Mycobacterium leprae* viability by polymerase chain reaction amplification of 71-kDa heat shock protein mRNA. *J. Infect.* 168, 799–800.
- Pitkaranta, M., Meklin, T., Hyvarinen, A., Nevalainen, A., Paulin, L., Auvinen, P., & Rintala, H. (2008). Analysis of fungal flora in indoor dust by ribosomal DNA sequence analysis, quantitative PCR, and culture. *Appl. Environ. Microbiology*. 74(1), 233-244
- Pitkaranta, M., Meklin, T., Hyvarinen, A., Nevalainen, A., Paulin, L., Auvinen, P., Lignell, U., & Rintala, H. (2011). Molecular profiling of fungal communities in moisture damaged buildings before and after remediation - a comparison of culture-dependent and culture-independent methods. *BMC Microbiology*. 11, 235.
- Portnoy, J. M., Kwak, K., Dowling, P., VanOsdol, T., & Barnes, C. (2005). Health effects of indoor fungi. *Annals of Allergy, Asthma & Immunology*, 94(3), 313-320.
- Pranoto, Y., Rakshit, S. K., & Salokhe, V. M. (2005). Enhancing antimicrobial activity of chitosan films by incorporating garlic oil, potassium sorbate and nisin. *LWT-Food Science and Technology*, 38(8), 859-865.

- Prussin, A. J., & Marr, L. C. (2015). Sources of airborne microorganisms in the built environment. *Microbiome*, 3(1), 78.
- Pryce-Miller, E., Aanensen, D., Vanittanakom, N., & Fisher, M. C. (2008). Environmental detection of *Penicillium marneffei* and growth in soil microcosms in competition with *Talaromyces stipitatus*. *Fungal Ecology*, 1(1), 49–56.
- Rani, P. (2015). The Impact of Adaptive Reusing Heritage Building as Assessed by the Indoor Air Quality Case study: UNESCO World Heritage Site Penang. *Procedia-Social and Behavioral Sciences*, 179, 297-307.
- Rawsthorne H, & Phister TG (2009) Detection of viable *Zygosaccharomyces bailii* in fruit juices using ethidium monoazide bromide and real-time PCR. *Int J Food Microbiol*, 131, 246 – 250.
- Robbins, C. A., Swenson, L. J., Nealley, M. L., Kelman, B. J., & Gots, R. E. (2000). Health effects of mycotoxins in indoor air: a critical review. *Applied occupational and environmental hygiene*, 15(10), 773-784.
- Ruddick, J. N. (2008). Biocide depletion: chemical, physical, and photodegradation. In *ACS symposium series* (Vol. 982, pp. 285-311). Oxford University Press.
- Ruiz-Capillas, C., & Nollet, L. M. (Eds.). (2015). *Flow Injection Analysis of Food Additives* (Vol. 1). CRC Press.
- Rys, P. N., & Persing, D. H. (1993). Preventing false positives: quantitative evaluation of three protocols for inactivation of polymerase chain reaction amplification products. *Journal of clinical microbiology*, 31(9), 2356-2360.
- Saari, A., Tissari, T., Valkama, E & Seppanen, O. (2005) The Effect of Redesigned Floor Plan, Occupant Density and the Quality of Indoor Climate on the Cost of Space, Productivity and Sick Leave in an Office Building – A Case Study. *Journal of Building and Environment*, 41(12), 1961-1972.

- Sambrook, J., Fritsch, E. F., & Maniatis, T. (1989). Quantitation of DNA and RNA. *Molecular cloning: a laboratory manual*, 3. New York: Cold Spring Harbor Laboratory Press.
- Sambrook, J. & Russell, D. W. (2006). Agarose gel electrophoresis. *Cold Spring Harbor Protocols*, 2006(1), pdb-prot4020.
- Samson, R. A., Houbroken, J. A. M. P., Kuijpers, A. F., Frank, J. M., & Frisvad, J. C. (2004). New ochratoxin A or sclerotium producing species in *Aspergillus* section *Nigri*. *Stud Mycol*, 50, 45-61.
- Samson, R. A., Houbroken, J., Thrane, U., Frisvad, J. C., & Andersen, B. (2010). Food and indoor fungi. *CBS laboratory manual series*.
- Samson, R. A., Yilmaz, N., Houbroken, J., Spierenburg, H., Seifert, K. A., Peterson, S. W., Varga, J. & Frisvad, J. C. (2011). Phylogeny and nomenclature of the genus *Talaromyces* and taxa accommodated in *Penicillium* subgenus *Biverticillium*. *Studies in Mycology*, 70(1), 159-183.
- Sánchez, G., Elizaquível, P., & Aznar, R. (2012). Discrimination of infectious hepatitis A viruses by propidium monoazide real-time RT-PCR. *Food and environmental virology*, 4(1), 21-25.
- Schweigkofler, W., O'Donnell, K., & Garbelotto, M. (2004). Detection and quantification of airborne conidia of *Fusarium circinatum*, the causal agent of pine pitch canker, from two California sites by using a real-time PCR approach combined with a simple spore trapping method. *Applied and Environmental Microbiology*, 70(6), 3512-3520.
- Sedlbauer, K. (2001). Prediction of mould fungus formation on the surface of and inside building components. *Fraunhofer Institute for Building Physics*.
- Sekhar, S. C., Tham, K. W., & Cheong, K. W. (2003). Indoor air quality and energy performance of air-conditioned office buildings in Singapore. *Indoor air*, 13(4), 315-331.

- Serpaggi, V., Remize, F., Recorbet, G., Gaudot-Dumas, E., Sequeira-Le Grand, A., & Alexandre, H. (2012). Characterization of the “viable but nonculturable” (VBNC) state in the wine spoilage yeast *Brettanomyces*. *Food microbiology*, 30(2), 438-447.
- Sheridan, G.E.C., Masters, C.I., Shallcross, J.A., & Mackey, B.M. (1998). Detection of mRNA by reverse transcription-PCR as an indicator of viability in *Escherichia coli* cells. *Appl. Environ. Microbiol.* 64, 1313–1318.
- Shi, H., Xu, W., Trinh, Q., Luo, Y., Liang, Z., Li, Y., & Huang, K. (2012). Establishment of a viable cell detection system for microorganisms in wine based on ethidium monoazide and quantitative PCR. *Food Control*.
- Shin, J. H., Nolte, F. S., Holloway, B. P., & Morrison., C. J. (1999). Rapid identification of up to three *Candida* species in a single reaction tube by a 59 exonuclease assay using fluorescent DNA probes. *J. Clin. Microbiol.* 37,165–170.
- Silva, D. M., Batista, L. R., Rezende, E. F., Fungaro, M. H. P., Sartori, D., & Alves, E. (2011). Identification of fungi of the genus *Aspergillus* section *nigri* using polyphasic taxonomy. *Brazilian Journal of Microbiology*, 42(2), 761-773.
- Silva, M. A., Iamanaka, B. T., Taniwaki, M. H., & Kieckbusch, T. G. (2013). Evaluation of the antimicrobial potential of alginate and alginate/chitosan films containing potassium sorbate and natamycin. *Packaging Technology and Science*, 26(8), 479-492.
- Sivasubramani SK, Niemeier RT, Reponen T, & Grinshpun SA. (2004) Assessment of the aerosolization potential for fungal spores in moldy homes. *Indoor Air*, 14, 405–412.

- Smilanick, J. L., Mansour, M. F., Gabler, F. M., & Sorenson, D. (2008). Control of citrus postharvest green mold and sour rot by potassium sorbate combined with heat and fungicides. *Postharvest Biology and Technology*, 47(2), 226-238.
- Smith, J. E., & Moss, M. O. (1985). *Mycotoxins. Formation, analysis and significance*. John Wiley & Sons Ltd..
- Soejima, T., Iida, K., Qin, T., Taniai, H., Seki, M., & Yoshida, S.I. (2008). Method to detect only live bacteria during PCR amplification. *J. Clin. Microbiol.* 46, 2305–2313.
- Spreadbury, C., Holden, D., Aufauvre-Brown, A., Bainbridge, B., & Cohen, J. (1993). Detection of *Aspergillus fumigatus* by polymerase chain reaction. *Journal of clinical microbiology*, 31(3), 615-621.
- Stryjawska-Sekulska, M., Piotraszewski-Pajak, A., Szyszka, A., Nowicki, M., & Filipiak, M. (2007). Microbiological quality of indoor air in university rooms. *Polish Journal of Environmental Studies*, 16(4), 623.
- Sudharsanam, S., Mathias, S., Barani, R., Sugumar, M., Janardhanan, J., Annamalai, R., Swaminathan, S., & Srikanth, P. (2015). Fate of airborne coagulase-negative staphylococci. *International Journal of Infection Control*, 11(4).
- Sullivan, D., Moran, G., & Coleman, D. (2005) in Kavanagh, K (Ed.), Fungi, Biology and Applications, *John Wiley & Sons*, England, 171-217.
- Susca, A., Stea, G., Mulè G., & Perrone, G. (2007). Polymerase chain reaction (PCR) identification of *Aspergillus niger* and *Aspergillus tubingensis* based on the calmodulin gene. *Food Additives and Contaminants*, 24(10), 1154–1160.
- Sutton, S. (2011). Accuracy of plate counts. *Journal of validation technology*, 17(3), 42. March 13, 2013. http://www.microbiol.org/wp-content/uploads/2010/07/Sutton.jvt_2011.17_3.pdf

- Tang, C. M., Holden, D. W., Aufauyre-Brown, A., & Cohen, J. (1993). The detection of *Aspergillus* spp. by the polymerase chain reaction and its evaluation in bronchoalveolar lavage fluid. *American Review of Respiratory Disease*, 148, 1313-1313.
- Tichenor, B. A., Sparks, L. A., White, J. B., & Jackson, M. D. (1990). Evaluating sources of indoor air pollution. *Journal of the Air & Waste Management Association*, 40(4), 487-492.
- Tsubouchi, H., Yamamoto, K., Hisada, K., Sakabe, Y., & Udagawa, S. I. (1987). Effect of roasting on ochratoxin A level in green coffee beans inoculated with *Aspergillus ochraceus*. *Mycopathologia*, 97(2), 111-115.
- Tulamait, A., Laghi, F., Mikrut, K., Carey, R. B., & Budinger, G. S. (2005). Potassium sorbate reduces gastric colonization in patients receiving mechanical ventilation. *Journal of critical care*, 20(3), 281-287.
- Türe, H., Gölstedt, M., & Hedenqvist, M. S. (2012). Antimicrobial compression-moulded wheat gluten films containing potassium sorbate. *Food Research International*, 45(1), 109-115.
- Udagawa, S.I., Tsubouchi, H., & Toyazaki, N. (1996). Isolation and identification of *Neosartorya* species from house dust as hazardous indoor pollutants. *Mycoscience*. 37, 217-222.
- Storey, E., Dangman, K. H., Schenck, P., DeBernardo, R. L., Yang, C. S., Bracker, A., & Hodgson, M. J. (2004). Guidance for clinicians on the recognition and management of health effects related to mold exposure and moisture indoors. Farmington, CT: University of Connecticut Health Center, Division of Occupational and Environmental Medicine. *Center for Indoor Environments and Health*.
- Van Burik, J. A., Schreckhise, R. W., White, T. C., Bowden, R. A., & Myerson, D. (1998). Comparison of six extraction techniques for isolation of DNA from filamentous fungi. *Medical mycology*, 36(5), 299-303.

- Varga, J., Kevei, E., Rinyu, E., T éren, J., & Kozakiewicz, Z. (1996). Ochratoxin production by *Aspergillus* species. *Applied and Environmental Microbiology*, 62(12), 4461-4464.
- Verma, R. K., Chaurasia, L., & Katiyar, S. (2008). Potential antifungal plants for controlling building fungi. *Natural Product Radiance*, 7(4), 374-387.
- Vesper, S., McKinstry, C., Hartmann, C., Neace, M., Yoder, S., & Vesper, A. (2008). Quantifying fungal viability in air and water samples using quantitative PCR after treatment with propidium monoazide (PMA). *Journal of microbiological methods*, 72(2), 180-184.
- Viegas, C., Almeida-Silva, M., Gomes, A. Q., Wolterbeek, H. T., & Almeida, S. M. (2014). Fungal contamination assessment in Portuguese elderly care centers. *Journal of Toxicology and Environmental Health, Part A*, 77(1-3), 14-23.
- Viviani, M. A., Hill, J. O., & Dixon, D. M. (1993). *Penicillium marneffei*: dimorphism and treatment. In *Dimorphic Fungi in Biology and Medicine* (pp. 413-422). Springer US.
- Wardah, A.R., Hafizi, R., Siti Nurdijati, B. & Salleh, B. (2011). Incidence and remediation of fungi in sick building in Malaysia- A case study. *Aerobiologia*. DOI: 10.1007/s10453-09226-y.
- Wensing, M., Uhde, E., & Salthammer, T. (2005). Plastics additives in the indoor environment—flame retardants and plasticizers. *Science of the Total Environment*, 339(1), 19-40.
- White, T. J., Bruns, T., Lee, S. J. W. T., & Taylor, J. L. (1990). Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. *PCR protocols: a guide to methods and applications*, 18(1), 315-322

- Williams, R. H., Ward, E., & McCartney, H. A. (2001). Methods for integrated air sampling and DNA analysis for detection of airborne fungal spores. *Applied and environmental microbiology*, 67(6), 2453-2459.
- World Health Organization. (2006). Almost a quarter of all disease caused by environmental exposure. In *Almost a quarter of all disease caused by environmental exposure*.
- Wolkoff, P., Wilkins, C. K., Clausen, P. A., & Larsen, K. (1993). Comparison of volatile organic compounds from processed paper and toners from office copiers and printers: methods, emission rates, and modeled concentrations. *Indoor air*, 3(2), 113-123.
- Wong, L.T., Mui, K.W., Hui, P.S., Chan, W.Y., & Law, A.K.Y. (2008). Thermal Environmental Interference with Airborne bacteria and Fungi Level in Air-Conditioning Offices, *Indoor and Built Environment*, 17(2), 122-127.
- Wu, Z., Wang, X.R. & Blomquist G. (2002). Evaluation of PCR primers and PCR conditions for specific detection of common airborne fungi. *J Environ Monit*, 4, 377-382.
- Wu, Z., Blomquist, G., Westermarck, S. O., & Wang, X. R. (2002). Application of PCR and probe hybridization techniques in detection of airborne fungal spores in environmental samples. *Journal of Environmental Monitoring*, 4(5), 673-678.
- Yamakami, Y., Hashimoto, A., Tokimatsu, I., & Nasu, M. (1996). PCR detection of DNA specific for *Aspergillus* species in serum of patients with invasive aspergillosis. *Journal of Clinical Microbiology*, 34(10), 2464-2468.
- Yañez, M.A., Nocker, A., Soria-Soria, E., Múrtula, R., Martinez, L. & Catalán, V., (2011). Quantification of viable *Legionella pneumophila* cells using propidium monoazide combined with quantitative PCR. *J. Microbiol. Methods* 85, 124-130.

- Yang, C.S. (2004). What do fungal PCR results mean? Emlab.com. Retrieved September 15, 2013, from <http://www.emlab.com/media/resources/Fungi-and-PCR.pdf>.
- Yap, J., Toh, Z. A., Goh, V., Ng, L. C., & Vesper, S. (2009). Assessment of mold concentrations in Singapore shopping centers using mold-specific quantitative PCR (MSQPCR) analysis. *Indian journal of microbiology*, 49(3), 290-293.
- Yassin, M.F. & Almouqatea, S. (2010). Assessment of Airborne bacteria and fungi in an indoor and outdoor environment. *Int. J. Environ. Sci. Tech.*, 7(3), 535-544.
- Yau, Y.H., Chew, B.T., & Saifullah, A.Z.A. (2012) Studies on the indoor air quality of Pharmaceutical Laboratories in Malaysia. *International Journal of Sustainable Built Environment*, 1, 110–124.
- Yilmaz, N., Houbraken, J., Hoekstra, E. S., Frisvad, J. C., Visagie, C. M., & Samson, R. A. (2012). Delimitation and characterisation of *Talaromyces purpurogenus* and related species. *Persoonia-Molecular Phylogeny and Evolution of Fungi*, 29(1), 39-54.
- Zhang, R. Z., Tsai, S. S., Chao, H. R., Shieh, Y. C., & Chuang, K. P. (2016). Innovation of a New Virus-Like Nanoparticle Vaccine System for the Specific Aerosol Relative Disease. *Aerosol and Air Quality Research*, 16(10), 2421-2427.
- Zielińska-Jankiewicz, K., Kozajda, A., Piotrowska, M., & Szadkowska-Stańczyk, I. (2008). Microbiological contamination with moulds in work environment in libraries and archive storage facilities. *Annals of Agricultural and Environmental Medicine*, 15(1).